

For Research Use Only

Phospho-mTOR (Ser2448) Recombinant antibody, PBS Only



Catalog Number: 80596-1-PBS

Basic Information

Catalog Number: 80596-1-PBS	GenBank Accession Number: BC117166	Purification Method: Protein A purification
Size: 100ug, Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 2475	CloneNo.: 3L18
Source: Rabbit	Full Name: FK506 binding protein 12-rapamycin associated protein 1	
Isotype: IgG	Calculated MW: 289 kDa	
	Observed MW: 250-289 kDa	

Applications

Tested Applications:
WB, IF, Indirect ELISA

Species Specificity:
Human, Rat

Background Information

mTOR, also named as FRAP1, FRAP, FRAP2 and RAPT1, belongs to the PI3/PI4-kinase family. mTOR is a Ser/Thr protein kinase that functions as an ATP and amino acid sensor to balance nutrient availability and cell growth. mTOR is kinase subunit of both mTORC1 and mTORC2, which regulate cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino-acids. mTORC2 is also activated by growth factors, but seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTOR is phosphorylated at Ser2448 via the PI3 kinase/Akt signaling pathway and autophosphorylated at Ser2481. mTOR plays a key role in cell growth and homeostasis and may be abnormally regulated in tumors.

Storage


Storage:
Store at -80°C.

Storage Buffer:
PBS Only


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Non-treated and Calyculin A treated HEK-293 cells were subjected to SDS PAGE followed by western blot with 80596-1-RR (Phospho-mTOR (Ser2448) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with RICTOR antibody (66867-2-Ig) subsequently. This data was developed using the same antibody clone with 80596-1-PBS in a different storage buffer formulation.



Non-treated and Calyculin A treated HeLa cells were subjected to SDS PAGE followed by western blot with 80596-1-RR (Phospho-mTOR (Ser2448) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with RICTOR antibody (66867-2-Ig) subsequently. This data was developed using the same antibody clone with 80596-1-PBS in a different storage buffer formulation.



Calyculin A treated HSC-T6 cells were subjected to SDS PAGE followed by western blot with 80596-1-RR (Phospho-mTOR (Ser2448) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80596-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed PMA treated HEK-293 cells using Phospho-mTOR (Ser2448) antibody (80596-1-RR, Clone: 3L18) at dilution of 1:1000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red). This data was developed using the same antibody clone with 80596-1-PBS in a different storage buffer formulation.