For Research Use Only

## Phospho-TSC2 (Thr1462) Recombinant antibody

Catalog Number:80698-1-RR



**Basic Information** 

Catalog Number: GenBank Accession Number:

80698-1-RR BC150300 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 7249

Nanodrop; **UNIPROT ID:** Source: P49815

Rabbit Full Name:

Isotype: tuberous sclerosis 2 IgG Calculated MW:

> 1807 aa, 201 kDa Observed MW: 200 kDa

**Applications** 

**Tested Applications:** 

WB, ELISA

Species Specificity:

Human

**Purification Method:** Protein A purification

CloneNo.: 1L20

Recommended Dilutions:

WB 1:2000-1:10000

Positive Controls:

WB: Calyculin A treated HEK-293 cells,

**Background Information** 

TSC2, also named Tuberin, plays an essential role in the cellular energy response pathway. TSC2 forms a physical and functional complex with TSC1 that can inhibit the nutrient-mediated or growth factor-stimulated phosphorylation of S6K1 and EIF4EBP1 by negatively regulating mTORC1 signaling. It has been reported that TSC2 can be phosphorylated at Thr1462 by Akt/PKB involved in stimulating cell growth and is activated by insulin. (PMID: 14651849, 12172553)

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

W: ptglab.com

## Selected Validation Data



Non-treated and Calyculin A treated HEK-293 cells were subjected to SDS PAGE followed by western blot with 80698-1-RR (Phospho-TSC2 (Thr1462) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with mTOR antibody (66888-1-lg) as the loading control.