For Research Use Only

TGFBI/BIGH3 Recombinant antibody

Catalog Number:80805-4-RR



Purification Method:

Protein A purification

Recommended Dilutions:

WB: 1:2000-1:10000

CloneNo.:

243108F7

Basic Information

Catalog Number: GenBank Accession Number:

100ul , Concentration: 1000 µg/ml by 7045
Nanodrop; ENSEMBL Gene ID:
Source: ENSG00000120708

Rabbit UNIPROT ID: Isotype: Q15582 IgG Full Name:

Immunogen Catalog Number: transforming growth factor, beta-EG1233 induced, 68kDa

Calculated MW: 75kDa

Observed MW: 68 kDa

Applications

Tested Applications: WB, ELISA

Species Specificity:

human

Positive Controls:

WB: HeLa cells, A549 cells

Background Information

TGFBI, also named as BIGH3, Kerato-epithelin and RGD-CAP, binds to type I, II, and IV collagens. TGFBI is an adhesion protein which may play an important role in cell-collagen interactions. In cartilage, it may be involved in endochondral bone formation. TGFBI is an extracellular matrix adaptor protein, it has been reported to be differentially expressed in transformed tissues. TGFBI is a predictive factor of the response to chemotherapy, and suggest the use of TGFBI-derived peptides as possible therapeutic adjuvants for the enhancement of responses to chemotherapy. (PMID:20509890) Defects in TGFBI are the cause of epithelial basement membrane corneal dystrophy (EBMD). Defects in TGFBI are the cause of corneal dystrophy Groenouw type 1 (CDGG1). Defects in TGFBI are the cause of corneal dystrophy lattice type 1 (CDL1). Defects in TGFBI are a cause of corneal dystrophy Thiel-Behnke type (CDTB). Defects in TGFBI are the cause of Reis-Buecklers corneal dystrophy (CDRB). Defects in TGFBI are the cause of lattice corneal dystrophy type 3A (CDL3A). Defects in TGFBI are the cause of Avellino corneal dystrophy (ACD).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 80805-4-RR (beta IG-H3/TGFBI antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.