

For Research Use Only

# HDAC3 Recombinant antibody

Catalog Number: 81211-1-RR



## Basic Information

Catalog Number:	81211-1-RR	GenBank Accession Number:	BC000614	Purification Method:	Protein A purification
Size:	100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI):	8841	CloneNo.:	2N1
Source:	Rabbit	UNIPROT ID:	O15379	Recommended Dilutions:	WB 1:5000-1:50000 IHC 1:50-1:500
Isotype:	IgG	Full Name:	histone deacetylase 3		
Immunogen Catalog Number:	AG0396	Calculated MW:	49 kDa		
		Observed MW:	49 kDa		

## Applications

Tested Applications:  
WB, IHC, ELISA

Species Specificity:  
Human, mouse, rat

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

Positive Controls:

**WB** : HeLa cells, HepG2 cells, MOLT-4 cells, Caco-2 cells, Jurkat cells, K-562 cells, NIH/3T3 cells, C6 cells

**IHC** : human thyroid cancer tissue, human liver cancer tissue

## Background Information

Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. Histone deacetylase (HDAC) and histone acetyltransferase (HAT) are enzymes that regulate transcription by selectively deacetylating or acetylating the (-amino groups of lysines located near the amino termini of core histone proteins. At least 4 classes of HDAC were identified. HDAC3 is a class I HDAC. HDAC3 has histone deacetylase activity and may participate in the regulation of transcription through its binding with the zinc-finger transcription factor YY1. HDAC3 can also down-regulate p53 function and thus modulate cell growth and apoptosis. The gene encoding HDAC3 is regarded as a potential tumor suppressor gene.

## Storage


Storage:  
Store at -20°C. Stable for one year after shipment.  
Storage Buffer:  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 81211-1-RR (HDAC3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human thyroid cancer tissue slide using 81211-1-RR (HDAC3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 81211-1-RR (HDAC3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 81211-1-RR (HDAC3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human thyroid cancer tissue slide using 81211-1-RR (HDAC3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 81211-1-RR against Human HDAC3 were performed. The affinity constant is 0.213 nM.