### For Research Use Only

# HA Tag Recombinant antibody

Catalog Number:81290-1-RR 44 Publications



**Basic Information** 

Catalog Number:

81290-1-RR

100ul, Concentration: 750 ug/ml by Nanodrop;

Source: Rabbit

Isotype:

GenBank Accession Number:

GeneID (NCBI):

6

Full Name: HA Tag Purification Method:

Protein A purification

CloneNo.:

6|21

Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IF/ICC 1:200-1:800

**Applications** 

Tested Applications: WB, IF/ICC, IP, ELISA Cited Applications:

WB, IF, IP, CoIP
Species Specificity:
recombinant protein

Cited Species: human

Positive Controls:

WB: Transfected HEK-293 cells,
IP: Transfected HEK-293T cells,
IF/ICC: Transfected HEK-293 cells,

## **Background Information**

Protein tags are a protein or peptide sequences located either on the C- or N- terminal of the target protein, which can facilitate solubility, detection, purification, localization, and expression of the target protein. The HA tag corresponds to amino acid residues YPYDVPDYA of a surface glycoprotein-human influenza virus hemagglutinin (HA). The HA tag is commonly used for a variety of research applications including chromatin immunoprecipitation, ELISA, flow cytometry, western blotting, immunocytochemistry/immunofluorescence among of others.

## Notable Publications

Author	Pubmed ID	Journal	Application
Zhengzheng Li	39912421	Adv Sci (Weinh)	WB
Linlin Xu	39893205	Cell Death Discov	IP
Xiannan Meng	39844192	J Transl Med	WB,IP

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3  $\,$ 


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


### **Selected Validation Data**




HA-tag recombinant protein, non-transfected HEK-293 cells, transfected HEK-293 cells were subjected to SDS PAGE followed by western blot with 81290-1-RR (HA Tag antibody) at dilution of 1:30000 incubated at room temperature for 1.5 hours.




IP result of anti-HA Tag (IP:81290-1-RR, 4ug; Detection:81290-1-RR 1:10000) with Transfected HEK-293T cells lysate 400 ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed Transfected HEK-293 cells using HA Tag antibody (81290-1-RR, Clone: 6121) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1x10^6 Transfected HEK-293 cells were intracellularly stained with 0.25 ug HA Tag Recombinant antibody (81290-1-RR, Clone:6l21) and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), and 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (4% PFA) fixed Transfected HEK-293 cells using HA Tag antibody (81290-1-RR, Clone: 6l21) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).