## For Research Use Only

## **VEGFA** Recombinant antibody

Catalog Number:81323-2-RR



**Purification Method:** 

**Basic Information** 

Catalog Number: GenBank Accession Number:

Protein A purification BC065522

GeneID (NCBI): CloneNo.: 100ul, Concentration: 1000 µg/ml by 7422 7H10

Nanodrop; Recommended Dilutions: Source: vascular endothelial growth factor A WB 1:5000-1:50000

Rabbit Calculated MW: Isotype: 412 aa, 46 kDa IgG Observed MW: 40 kDa

Immunogen Catalog Number:

AG13500

81323-2-RR

**Applications Tested Applications:** Positive Controls:

WB, ELISA WB: MCF-7 cells, T-47D cells, A431 cells, MDA-MB-231

Species Specificity: cells, HUVEC cells, RAW 264.7 cells

Human, mouse

## **Background Information**

VEGFA, also named as VEGF or VPF, belongs to the PDGF/VEGF growth factor family. It is a growth factor active in angiogenesis, vasculogenesis and endothelial cell growth. VEGFA induces endothelial cell proliferation, promotes cell migration, inhibits apoptosis and induces permeabilization of blood vessels. It binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin. Defects in VEGFA are associated with microvascular complications of diabetes type 1 (MVCD1). VEGFA has 17 isoforms with MW from 16 to 45 kDa. Some isoforms have homodimer forms (e.g.; VEGFA189 38 kDa or VEFGA110 34 kDa). VEGF-A exists in at least seven homodimeric isoforms. The monomers consist of 121, 145, 148, 165, 183, 189, or 206 amino acids (PMID:15602010).

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 81323-2-RR (VEGFA antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.