

For Research Use Only

FTO Recombinant antibody

Catalog Number: 81471-1-RR



Basic Information

Catalog Number: 81471-1-RR	GenBank Accession Number: NM_001080432	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 79068	CloneNo.: 4L4
Source: Rabbit	Full Name: fat mass and obesity associated	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:500-1:2000 IF 1:50-1:500
Isotype: IgG	Calculated MW: 58 kDa	
Immunogen Catalog Number: AG26095	Observed MW: 58 kDa	

Applications

Tested Applications:

IF, IHC, WB, ELISA

Species Specificity:

Human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HeLa cells, HEK-293T cells, LNCaP cells, Jurkat cells

IHC : human liver cancer tissue,

IF : HepG2 cells,

Background Information

Fat mass and obesity-associated protein (FTO) has efficient oxidative demethylation activity targeting the abundant N6-methyladenosine (m6A) residues in RNA in vitro. Variants in the FTO (fat mass and obesity associated) gene are associated with increased body mass index in humans.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 81471-1-RR (FTO antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 81471-1-RR (FTO antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using FTO antibody (81471-1-RR, Clone: 4L4) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).