For Research Use Only

DMT1 Recombinant antibody

Catalog Number:81609-1-RR



Basic Information

Catalog Number: GenBank Accession Number:

 81609-1-RR
 BC002592

 Size:
 GeneID (NCBI):

100ul, Concentration: 1000 µg/ml by 4891

anodrop; Full Name:

Source: solute carrier family 11 (proton-Rabbit coupled divalent metal ion Isotype: transporters), member 2

IgGCalculated MW:Immunogen Catalog Number:568 aa, 62 kDaAG14314Observed MW:

70 kDa

Applications

Tested Applications:

IP, WB, ELISA

Species Specificity:
Human, mouse, rat

Positive Controls:

WB: HeLa cells, HepG2 cells, HEK-293 cells, HuH-7 cells, Caco-2 cells, Neuro-2a cells, HSC-T6 cells

Purification Method:

CloneNo.:

1113

for WB

Protein A purification

Recommended Dilutions:

IP 0.5-4.0 ug for IP and 1:1000-1:4000

WB 1:5000-1:50000

IP: HEK-293 cells.

Background Information

SLC11A2 (also known as DMT1, Nramp2, and DCT1) is a member of the divalent cation transporters that plays a central role in iron homeostasis. SLC11A2 is widely expressed in many tissues including brain, kidney, testis, duodenum and placenta. As a membrane protein, SLC11A2 is localized on the apical membrane of enterocytes as well as in transferrin-cycle endosomes. Four isoforms of SLC11A2 exist due to the alternative splicing. They differ at the NH2 and COOH termini but share a common central domain. A variety of molecular weights of SLC11A2 in western blot analysis, ranging from 50 to 100 kDa, has been reported in different cells and species. The differences in molecular weights may be attributed to the level of glycosylation, proteolysis, or the membrane protein itself. This antibody detected various forms of SLC11A2 of 45-100 kDa.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 81609-1-RR (DMT1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



IP result of anti-DMT1(IP:81609-1-RR, 4ug; Detection:81609-1-RR 1:2000) with HEK-293 cells lysate 1480 ug.