

For Research Use Only

# Hsc70 Recombinant antibody

Catalog Number: 81719-2-RR

Featured Product



## Basic Information

Catalog Number:

81719-2-RR

Size:

100ul, Concentration: 500 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1018

GenBank Accession Number:

BC007276

GeneID (NCBI):

3312

UNIPROT ID:

P11142

Full Name:

heat shock 70kDa protein 8

Calculated MW:

70 kDa

Observed MW:

70 kDa

Purification Method:

Protein A purification

CloneNo.:

3H9

Recommended Dilutions:

WB 1:5000-1:50000

## Applications

Tested Applications:

WB, ELISA

Species Specificity:

Human, mouse, rat

Positive Controls:

WB : HeLa cells, HepG2 cells, Jurkat cells, NIH/3T3 cells, MCF-7 cells, PC-12 cells

## Background Information

HSPA8 (also known as HSC70) is a member of the HSPA (HSP70) family of heat-shock proteins which are highly conserved chaperons implicated in protein folding, protein refolding, protein transport, and protein targeting. HSPA8 is a constitutively expressed cytosol/nuclear protein able to translocate between cytoplasm and nucleus. Recently it has been reported that HSPA8 can interact with  $\alpha$ -synuclein, the critical pathological protein of Parkinson's disease, indicating its implication in neurodegenerative disease. (PMID: 21832061)

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 81719-2-RR (Hsc70 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



WB result of Hsc70 antibody (81719-2-RR; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Hsc70 transfected HeLa cells.