

For Research Use Only

Phospho-TAK1 (Thr187) Recombinant antibody

Catalog Number: 81785-1-RR



Basic Information

Catalog Number: 81785-1-RR	GenBank Accession Number: BC017715	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 6885	CloneNo.: 206
Source: Rabbit	UNIPROT ID: O43318	Recommended Dilutions: WB 1:20000-1:100000
Isotype: IgG	Full Name: mitogen-activated protein kinase kinase kinase 7	
	Calculated MW: 579 aa, 64 kDa	
	Observed MW: 75-85 kDa	

Applications

Tested Applications: WB, FC (Intra), ELISA	Positive Controls: WB : Calyculin A treated HEK-293 cells, Calyculin A treated HeLa cells
Species Specificity: human	

Background Information

MAP3K7(Mitogen-activated protein kinase kinase kinase 7) is also named TAK1 and belongs to the MAP kinase kinase kinase subfamily. It plays an important role in the cascades of cellular responses evoked by changes in the environment. It has been linked to interleukin-1 receptor and tumor necrosis factor receptor signaling (PMID: 16186825). It has 4 isoforms (53-55 kDa; 64-70 kDa and 75-80 kDa) produced by alternative splicing.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated and Calyculin A treated HEK-293 cells were subjected to SDS PAGE followed by western blot with 81785-1-RR (Phospho-TAK1 (Thr187) antibody) at dilution of 1:100000 incubated at room temperature for 1 hour. The membrane was stripped and re-blotted with Alpha Tubulin antibody as loading control.



1×10^6 HEK-293 cells untreated (dashed lines) or treated with Calyculin A were intracellularly stained with 0.13 μ g Phospho-TAK1 (Thr187) Recombinant antibody (81785-1-RR, Clone:206) and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.13 μ g Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.