For Research Use Only

USP9X Recombinant antibody, PBS Only

Catalog Number:81892-1-PBS Featured Product



Basic Information

Catalog Number:

81892-1-PBS

GenBank Accession Number: NM_001039590

Purification Method: Protein A purification

IgG

GeneID (NCBI): 100ug, Concentration: 1mg/ml by

CloneNo.: 1B23

Nanodrop:

Source: Rabbit

Q93008 Full Name:

UNIPROT ID:

Isotype:

ubiquitin specific peptidase 9, X-

linked

Calculated MW: 292 kDa Observed MW: 260-290 kDa

Applications

Tested Applications:

WB, IF/ICC, IP, Indirect ELISA

Species Specificity:

human, mouse, rat

Background Information


USP9X, also named as DFFRX, FAM and USP9, belongs to the peptidase C19 family. It may function as a ubiquitin $protein\ or\ polyubiquit in\ hydrolase\ involved\ both\ in\ the\ processing\ of\ ubiquit in\ precursors\ and\ of\ ubiquit in\ attack and\ of\ ubiquit in\ precursors\ and\ of\ ubiquit in\ attack and\ of\ ubiquit in\ precursors\ and\ of\ ubiquit in\ attack and\ of\ ubiquit in\ ubiquit in\ ubiquit in\ ubiquit in\ ubiquit in\ ubiquit in\$ proteins. It plays an important role regulatory role at the level of protein turnover by preventing degradation of proteins through the removal of conjugated ubiquitin. USP9X is an essential component of TGF-beta/BMP signaling cascade. Deubiqitination of SMAD4 by USP9X restores its competence to mediate TGF-beta signaling. USP9X regulates chromosome alignment and segregation in mitosis by regulating the localization of BIRC5/survivin to mitotic centromeres. The antibody is specific to USP9X. It has no cross reaction to USP9Y.

Storage

Storage: Store at -80°C. Storage Buffer: PBS Only

in USA), or 1(312) 455-8498 (outside USA)


Selected Validation Data




HEK-293 cells were subjected to SDS PAGE followed by western blot with 81892-1-RR (USP9X antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 81892-1-PBS in a different storage buffer formulation.




WB result of USP9X antibody (81892-1-RR; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-USP9X transfected HEK-293 cells. This data was developed using the same antibody clone with 81892-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using USP9X antibody (81892-1-RR, Clone: 1823) at dilution of 1:200 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 81892-1-PBS in a different storage buffer formulation.



IP result of anti-USP9X (IP:81892-1-RR, 4ug; Detection:81892-1-RR 1:3000) with HEK-293 cells lysate 1640 ug. This data was developed using the same antibody clone with 81892-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse testis tissue using USP9X antibody (81892-1-RR, Clone: 1B23) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 81892-1-PBS in a different storage buffer formulation.