

For Research Use Only

STX17 Recombinant antibody

Catalog Number: 81899-1-RR



Basic Information

Catalog Number: 81899-1-RR	GenBank Accession Number: BC111009	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 55014	CloneNo.: 4G3
Source: Rabbit	Full Name: syntaxin 17	Recommended Dilutions: WB 1:5000-1:50000 IF 1:200-1:800
Isotype: IgG	Calculated MW: 302 aa, 33 kDa	
Immunogen Catalog Number: AG12134	Observed MW: 33-35 kD	

Applications

Tested Applications:

IF, WB, ELISA

Species Specificity:

Human, Mouse, Rat

Positive Controls:

WB : human testis tissue, A375 cells, HEK-293 cells, HepG2 cells, L02 cells, mouse testis tissue, rat testis tissue

IF : HepG2 cells,

Background Information

SNAREs, soluble N-ethylmaleimide-sensitive factor-attachment protein receptors, are essential proteins for fusion of cellular membranes. Syntaxin 17 (STX17) is a SNARE of the autophagosome involved in autophagy through the direct control of autophagosome membrane fusion with the lysosome membrane. STX17 may also play a role in the early secretory pathway where it may maintain the architecture of the endoplasmic reticulum-Golgi intermediate compartment/ERGIC and Golgi and/or regulate transport between the endoplasmic reticulum, the ERGIC and the Golgi.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 81899-1-RR (STX17 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using STX17 antibody (81899-1-RR, Clone: 4G3) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).