For Research Use Only

Angiopoietin 1 Recombinant antibody, PBS Only

Catalog Number:81990-4-PBS



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

81990-4-PBS

GeneID (NCBI):

BC152411

Protein A purfication

100ug, Concentration: 1 mg/ml by

CloneNo.: 240764D10

Nanodrop:

UNIPROT ID: Q15389

Rabbit Isotype:

Full Name: angiopoietin 1

IgG Immunogen Catalog Number: Calculated MW: 498 aa, 58 kDa

AG18829

Observed MW: 70 kDa

Applications

Tested Applications:

WB, IHC, FC (Intra), ELISA

Species Specificity:

human


Background Information

Angiopoietin 1 is a 70-kDa secreted glycoprotein generated from vascular mural cells, pericytes, and certain other cells. Angiopoietin 1 participates in vascular differentiation through angiogenesis, which is the process of the growth and remodelling of existing vessels. Angiopoietin 1 is also involved in the maintenance and turnover of blood vessels in mature animals. Angiopoietin 1 binds to and activates the TEK/TIE2 receptor by inducing its $dimerization\ and\ tyrosine\ phosphorylation,\ playing\ an\ important\ role\ in\ the\ regulation\ of\ angiogenesis.$ Overexpression of Angiopoietin 1 has been proven to occur in malignant glioblastoma, neuroblastoma, non-small cell lung cancer, and other tumors.


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 81990-4-RR (Angiopoietin 1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 81990-4-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 81990-4-RR (Angiopoietin 1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 81990-4-PBS in a different storage buffer formulation.



1x10^6 HepG2 cells were intracellularly stained with 0.25 ug Angiopoietin 1 Recombinant antibody (81990-4-RR, Clone:240764D10) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 81990-4-PBS in a different storage buffer formulation.



Biolayer interferometry (BLL) kinetic assays of 81990-4-RR against Human ANGPT1 were performed. The affinity constant is 6.37 nM.