For Research Use Only

G3BP2 Recombinant antibody

Catalog Number:82080-6-RR



Basic Information

Catalog Number: GenBank Accession Number:

82080-6-RR BC011731 GeneID (NCBI):

100ul , Concentration: 1000 $\mu g/ml$ by 9908 Nanodrop: **UNIPROT ID:** Q9UN86 Rabbit Full Name:

Isotype: GTPase activating protein (SH3 domain) binding protein 2 IgG

Immunogen Catalog Number: Calculated MW:

482aa,54 kDa; 449aa,51 kDa AG9355

> Observed MW: 65-70 kDa

Applications

Tested Applications:

FC, IF, WB, ELISA Species Specificity: Human, mouse

Positive Controls:

WB: HeLa cells, K-562 cells, HEK-293 cells, A549 cells, Jurkat cells, MCF-7 cells, mouse cerebellum tissue

Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:2000-1:10000

IF 1:500-1:2000

CloneNo.:

230275G3

IF: sodium arsenite treated HeLa cells.

Background Information

Stress granules (SGs) are cytoplasmic mRNA-protein condensates formed in response to cellular stressors, such as oxidative stress, ultraviolet radiation, and viral infection (1). The Ras-GTPase-activating protein-binding proteins (G3BPs), consisting of G3BP1 and G3BP2, are key nucleating factors essential for SG formation. They function to the state of the statprotect RNAs from harmful conditions. G3BP2 is mainly distributed in the cytoplasm and participates in the formation of stress granules, cell differentiation, proliferation, and signal transduction. Accumulating evidence has demonstrated that aberrant expression of G3BP2 contributes to cancer initiation and progression, such as high expression of G3BP2 increasing cell stemness, metastasis and chemoresistance in breast cancer.

Storage

Store at -20°C. Stable for one year after shipment.


Aliquoting is unnecessary for -20°C storage

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82080-6-RR (G3BP2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed sodium arsenite treated HeLa cells using G3BP2 antibody (82080-6-RR, Clone: 230275G3) at dilution of 1:1000 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).



1x10^6 U2OS cells were intracellularly stained with 0.25 ug Anti-Human G3BP2 (82080-6-RR, Clone:230275G3) and APC-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L)(red), or 0.25 ug Rabbit 1gG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).