For Research Use Only

TCTN2 Recombinant antibody

Catalog Number:82121-1-RR



Basic Information

Catalog Number: GenBank Accession Number:

 82121-1-RR
 BC009112
 Protein A p

 Size:
 GeneID (NCBI):
 CloneNo.:

 100ul , Concentration: 1000 μg/ml by 79867
 1E17

Nanodrop; Full Name:

Source: tectonic family member 2

Rabbit Calculated MW:
Isotype: 697 aa, 77 kDa
IgG Observed MW:
Immunogen Catalog Number: 90 kDa

ΔG10725

Applications

Tested Applications:

WB, ELISA

Species Specificity:

Human, rat

Purification Method: Protein A purification

Recommended Dilutions:

WB: HEK-293 cells, A431 cells, HeLa cells, Jurkat cells,

WB 1:5000-1:50000

Background Information

TCTN2 (tectonic-2 or TECT2) is a type I membrane protein that belongs to the tectonic family, which consists of TCTN1, TCTN2 and TCTN3. Studies in mice suggest that tectonic may be involved in Hedgehog (Hh) signaling, and essential for ciliogenesis. Tectonic is component of the tectonic-like complex, a complex localized at the transition zone of primary cilia and acting as a barrier that prevents diffusion of transmembrane proteins between the cilia and plasma membranes. It exists two spilcing isoforms. The first aberrant transcript introduces 104 base pair from 13 and would delete 196 original amino acid, introduce two novel amino acids, and prematurely truncate the 697aa protein at residue number 504. The second transcript lacks exon 14 and would delete 195 original amino acids, introduce four novel amino acids, and prematurely truncate the 697 aa protein at residue number 507. The molecular weight of two isoforms are about 50kDa. We detected the 50kDa and 90kDa protein by immunoprecipitaion, but 50kDa protein may be the splicing isoform or non-specific band.

Positive Controls:

K-562 cells, RAW 264.7 cells

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data

Various lysates were subjected to SDS PAGE followed by western blot with 82121-1-RR (TCTN2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.