For Research Use Only

SIRT3 Recombinant antibody

Catalog Number:82210-1-RR



Basic Information

Catalog Number: GenBank Accession Number: 82210-1-RR BC001042

82210-1-RR BC001042 Protein A purification
Size: GeneID (NCBI): CloneNo.:

100ul , Concentration: 500 ug/ml by 23410 2P6
Nanodrop:

 Nanodrop;
 UNIPROT ID:
 Recommended Dilutions:

 Source:
 Q9NTG7
 WB 1:5000-1:50000

 Rabbit
 Full Name:
 IHC 1:200-1:800

Isotype: sirtuin (silent mating type

IgG information regulation 2 homolog) 3

Immunogen Catalog Number: (S. cerevisiae)
AG26280 Calculated MW:

44 kDa Observed MW: 28-30 kDa

Applications

Tested Applications: WB, IHC, ELISA

Species Specificity: human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HeLa cells, mouse heart tissue, rat heart tissue, HepG2 cells, HT-29 cells, mouse liver tissue, pig liver

Purification Method:

tissue, rat liver tissue

IHC: human prostate cancer tissue,

Background Information

Sirtuins are NAD+-dependent histone/protein deacetylases (HDAC) and SIRT3 is the only sirtuin whose increased expression has been shown to correlate with an extended life span in humans. It is localized in the mitochondrial matrix, where it regulates the acetylation levels of metabolic enzymes, including acetyl coenzyme A synthetase 2. SIRT3 is stress-responsive and its increased expression protects myocytes from genotoxic and oxidative stress-mediated cell death. full-length hSIRT3 (44 kDa) is an inert protein and that it is activated inside the mitochondria following deletion of 142 amino acids of the N-terminal segment. This processed form of SIRT3 is approximately 28 kDa and possesses deacetylase activity. The full SIRT3 is a 44 kDa protein and is processed by mitochondrial processing peptidase (MPP) to give a 28 kDa product.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82210-1-RR (SIRT3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 82210-1-RR (SIRT3 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 82210-1-RR (SIRT3 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLI) kinetic assays of 82210-1-RR against Human SIRT3 were performed. The affinity constant is 0.261 nM.