

For Research Use Only

# Phospho-NF- $\kappa$ B p65 (Ser468) Recombinant antibody



Catalog Number: 82335-1-RR **26 Publications**

## Basic Information

<b>Catalog Number:</b> 82335-1-RR	<b>GenBank Accession Number:</b> BC011603	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 100ul , Concentration: 500 $\mu$ g/ml by Nanodrop;	<b>GeneID (NCBI):</b> 5970	<b>CloneNo.:</b> 6N1
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> Q04206	<b>Recommended Dilutions:</b> WB 1:2000-1:10000
<b>Isotype:</b> IgG	<b>Full Name:</b> v-rel reticuloendotheliosis viral oncogene homolog A (avian)	
	<b>Calculated MW:</b> 65 kDa	
	<b>Observed MW:</b> 75 kDa	

## Applications

<b>Tested Applications:</b> WB, ELISA	<b>Positive Controls:</b> WB : Calyculin A treated HeLa cells, Calyculin A treated NIH/3T3 cells
<b>Cited Applications:</b> WB, IF	
<b>Species Specificity:</b> Human, Mouse	
<b>Cited Species:</b> human, rat, mouse, pig	

## Background Information

Nuclear factor  $\kappa$ B (NF- $\kappa$ B) is a collective term for a small family of dimeric transcription factors [comprising p65 (RelA) and RelB, c-Rel, p50/p105 (NF- $\kappa$ B1), and p52/p100 (NF- $\kappa$ B2)]. All NF- $\kappa$ B proteins share a Rel homology domain (RHD), which is responsible for DNA binding and dimerization. Only p65, RelB, and c-Rel contain potent transactivation domains within sequences from the C-terminal to the RHD. Exterior signals lead to the phosphorylation and degradation of the inhibitory complex I $\kappa$ B, which is modulated by the I $\kappa$ B kinase (IKK), and its degradation allows for the release of the typical NF- $\kappa$ B heterodimer, p65/p50, to translocate into the nucleus. NF- $\kappa$ B binds to its cognate DNA elements and can transcriptionally activate different target genes among which 200-500 genes have been implicated in cell survival/apoptosis, cell growth, immune response, and inflammation.

## Notable Publications

Author	Pubmed ID	Journal	Application
Yu Wang	38467242	Cell Signal	WB
Rumin Huang	38437888	J Ethnopharmacol	WB
Pengfei Dong	38429764	Eur J Med Res	WB

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Non-treated and Calyculin A treated various cells were subjected to SDS PAGE followed by western blot with 82335-1-RR (Phospho-NF-κB p65 (Ser468) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with Alpha Tubulin antibody (66031-1-Ig) and NF-κB p65 antibody (80979-1-RR) subsequently.