For Research Use Only

Cytokeratin 20 Recombinant antibody, PBS Only

www.ptglab.com

Catalog Number:82428-1-PBS

Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Protein A purification

82428-1-PBS

GeneID (NCBI):

CloneNo.: 2H10

Nanodrop:

100ug, Concentration: 1 mg/ml by

54474

BC031559

UNIPROT ID: P35900

Rabbit Isotype:

Full Name: keratin 20

Immunogen Catalog Number:

Calculated MW: 424 aa, 48 kDa

AG11308

IgG

Observed MW:

46 kDa

Applications

Tested Applications:

WB, IF, FC, IHC, Indirect ELISA

Species Specificity:

Human, mouse, rat, pig

Background Information


Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells, which are classified into two major sequence types. Type I keratins are a group of acidic intermediate filament proteins, including K9-K23, and the hair keratins Ha1-Ha8. Type II keratins are the basic or neutral courterparts to the acidic type I keratins, including K1-K8, and the hair keratins, Hb1-Hb6. Keratin 20 is a type I cytokeratin. It is a major cellular protein of mature enterocytes and goblet cells and is specifically found in the gastric and intestinal mucosa.

Storage


Storage: Store at -80°C. Storage Buffer:

PBS Only


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82428-1-RR (Cytokeratin 20 antibody) at dilution of 1:22000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 82428-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human appendicitis tissue slide using 82428-1-RR (Cytokeratin 20 antibody) at dilution of 1:1100 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 82428-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human appendicitis tissue slide using 82428-1-RR (Cytokeratin 20 antibody) at dilution of 1:1100 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 82428-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HT-29 cells using Cytokeratin 20 antibody (82428-1-RR, Clone: 2H10) at dilution of 1:1000 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 82428-1-PBS in a different storage buffer formulation.



1X10^6 HT-29 cells were intracellularly stained with 0.4 ug Anti-Human Cytokeratin 20 (82428-1-RR, Clone:2H10) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP, Clone:) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 82428-1-PBS in a different storage buffer