For Research Use Only

EIF3D Recombinant antibody

Catalog Number:82512-1-RR **Featured Product**



Basic Information

Catalog Number: GenBank Accession Number:

82512-1-RR BC000328 GeneID (NCBI): Size: CloneNo.: 100ul, Concentration: 1000 ug/ml by 8664 308

Nanodrop: **UNIPROT ID:** Recommended Dilutions: 015371 WB 1:5000-1:50000

Rabbit Full Name:

Isotype: eukaryotic translation initiation factor 3, subunit D IgG

Immunogen Catalog Number: Calculated MW: AG0268 66 kDa

Observed MW: 65 kDa

Purification Method: Protein A purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:1000-1:4000 IF/ICC 1:500-1:2000

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA

Species Specificity:

buffer pH 6.0

human, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

Positive Controls:

WB: HeLa cells, rat liver tissue, HEK-293 cells, HepG2

cells, A549 cells, A431 cells, Jurkat cells

IP: A549 cells,

IHC: human breast cancer tissue,

IF/ICC: U2OS cells,

Background Information

The mammalian translation initiation factor 3 (eIF3), is a multiprotein complex of ~600 kDa that binds to the 40 S ribosome and promotes the binding of methionyl-tRNAi and mRNA. The EIF3S7(p66) is the major RNA binding $subunit\ in\ this\ complex.\ Human\ el\ F3-p66\ shares\ 64\%\ sequence\ identity\ with\ a\ hypothetical\ Caenorhabditis\ elegans$ protein, presumably the p66 homolog. Deletion analyses of recombinant derivatives of eIF3-p66 show that the RNAbinding domain lies within an N-terminal 71-amino acid region rich in lysine and arginine.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com


Selected Validation Data




WB result of EIF3D antibody (82512-1-RR; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-EIF3D transfected HEK-293 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 82512-1-RR (EIF3D antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




rat liver tissue were subjected to SDS PAGE followed by western blot with 82512-1-RR (EIF3D antibody) at dilution of 1:30300 incubated at room temperature for 1.5 hours.




IP result of anti-EIF3D (IP:82512-1-RR, 4ug; Detection:82512-1-RR 1:5000) with A549 cells lysate 920 ug.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 82512-1-RR (EIF3D antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Sodium Citrate buffer (pH 6.0).




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 82512-1-RR (EIF3D antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Sodium Citrate buffer (pH 6.0).



Immunofluorescent analysis of (-20°C Methanol) fixed U2OS cells using EIF3D antibody (82512-1-RR, Clone: 308) at dilution of 1:1000 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol) fixed U2OS cells using EIF3D antibody (82512-1-RR, Clone: 3O8) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).



Biolayer interferometry (BLL) kinetic assays of 82512-1-RR against Human EIF3D were performed. The affinity constant is 5.83 nM.