For Research Use Only

## Cleaved Caspase 3/P17/P19 Recombinant antibody Catalog Number:82707-13-RR 3 Publications




Basic Information	Catalog Number: 82707-13-RR	GenBank Accession Number: NM_032991	Purification Method: Protein A purfication	
	Size:	GenelD (NCBI):	CloneNo.:	
	100ul , Concentration: 1000 ug/ml by		230527B2	
	Nanodrop;	UNIPROT ID:	Recommended Dilutions:	
	Source:	P42574	WB 1:5000-1:50000	
	Rabbit	Full Name:		
	Isotype: IgG	caspase 3, apoptosis-related cysteine peptidase		
		Calculated MW: 32 kDa		
		Observed MW: 17 kDa, 19 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, ELISA	WB : Staurosporine treated Jurkat cells,		
	Cited Applications: WB, IF			
	Species Specificity: human, mouse			
	Cited Species:			
	mouse			
Background Information	Caspases, a family of endoproteases, cell death. Initiator caspases (caspase (caspase-3, -6, and -7), which in turn e role in the activation of sterol regular leucine zipper domain and the memb proteins. This antibody can recognize	e-2, -8, -9, -10, -11, and -12) cleave execute apoptosis by cleaving targ tory element binding proteins (SRE orane attachment domain. Caspase p17 and p19 fragment of Caspase l length caspase-3. The cleaved P1	ory networks controlling inflammation and and activate downstream effector caspases eted cellular proteins. Caspase 3 plays a key EBPs) between the basic helix-loop-helix 2 Gan form heterocomplex with other 3. This antibody is specific for cleaved 7 and P19 fragment might form complex an	
	Caspases, a family of endoproteases, cell death. Initiator caspases (caspase (caspase-3, -6, and -7), which in turn e role in the activation of sterol regular leucine zipper domain and the memb proteins. This antibody can recognize caspase 3, and does not recognize ful shows at around 30-35 kDa by wester	e-2, -8, -9, -10, -11, and -12) cleave execute apoptosis by cleaving targ tory element binding proteins (SRE orane attachment domain. Caspase p17 and p19 fragment of Caspase l length caspase-3. The cleaved P1	and activate downstream effector caspases eted cellular proteins. Caspase 3 plays a key BPs) between the basic helix-loop-helix 3 can form heterocomplex with other 3. This antibody is specific for cleaved 7 and P19 fragment might form complex an	
	Caspases, a family of endoproteases, cell death. Initiator caspases (caspase (caspase-3, -6, and -7), which in turn e role in the activation of sterol regular leucine zipper domain and the memb proteins. This antibody can recognize caspase 3, and does not recognize ful shows at around 30-35 kDa by wester	e-2, -8, -9, -10, -11, and -12) cleave execute apoptosis by cleaving targ tory element binding proteins (SRE prane attachment domain. Caspase p17 and p19 fragment of Caspase I length caspase-3. The cleaved P1 n blot (PMID: 25501826).	and activate downstream effector caspases eted cellular proteins. Caspase 3 plays a key BPs) between the basic helix-loop-helix 3 can form heterocomplex with other 3. This antibody is specific for cleaved	
Background Information	Caspases, a family of endoproteases, cell death. Initiator caspases (caspase (caspase-3, -6, and -7), which in turn er role in the activation of sterol regular leucine zipper domain and the memb proteins. This antibody can recognize caspase 3, and does not recognize ful shows at around 30-35 kDa by wester Author Pul Fei Hu 39	e-2, -8, -9, -10, -11, and -12) cleave execute apoptosis by cleaving targ tory element binding proteins (SRE rane attachment domain. Caspase p17 and p19 fragment of Caspase l length caspase-3. The cleaved P1 n blot (PMID: 25501826).	and activate downstream effector caspases eted cellular proteins. Caspase 3 plays a key BPS) between the basic helix-loop-helix 3 can form heterocomplex with other 3. This antibody is specific for cleaved 7 and P19 fragment might form complex an Application WB	
	Caspases, a family of endoproteases, cell death. Initiator caspases (caspase (caspase-3, -6, and -7), which in turn er role in the activation of sterol regular leucine zipper domain and the memb proteins. This antibody can recognize caspase 3, and does not recognize ful shows at around 30-35 kDa by wester Author Pul Fei Hu 39' Bin Xia 39'	e-2, -8, -9, -10, -11, and -12) cleave execute apoptosis by cleaving targ tory element binding proteins (SRE rane attachment domain. Caspase p17 and p19 fragment of Caspase I length caspase-3. The cleaved P1 n blot (PMID: 25501826). bmed ID Journal 939534 J Membr Biol	and activate downstream effector caspases eted cellular proteins. Caspase 3 plays a key EBPs) between the basic helix-loop-helix 3 can form heterocomplex with other 3. This antibody is specific for cleaved 7 and P19 fragment might form complex and <b>Application</b> WB ophys WB,IF	
	Caspases, a family of endoproteases, cell death. Initiator caspases (caspase (caspase-3, -6, and -7), which in turn er role in the activation of sterol regular leucine zipper domain and the memb proteins. This antibody can recognize caspase 3, and does not recognize ful shows at around 30-35 kDa by wester Author Pul Fei Hu 39' Bin Xia 39'	e-2, -8, -9, -10, -11, and -12) cleave execute apoptosis by cleaving targ tory element binding proteins (SRE prane attachment domain. Caspase p17 and p19 fragment of Caspase l length caspase-3. The cleaved P1 n blot (PMID: 25501826). bmed ID Journal 939534 J Membr Biol 798643 Arch Biochem Bio	and activate downstream effector caspases eted cellular proteins. Caspase 3 plays a key BPs) between the basic helix-loop-helix 3 can form heterocomplex with other 3. This antibody is specific for cleaved 7 and P19 fragment might form complex an Application WB ophys WB,IF	
Notable Publications	Caspases, a family of endoproteases, cell death. Initiator caspases (caspase (caspase-3, -6, and -7), which in turn er role in the activation of sterol regular leucine zipper domain and the memb proteins. This antibody can recognize caspase 3, and does not recognize ful shows at around 30-35 kDa by wester Author Pul Fei Hu 39' Bin Xia 39'	e-2, -8, -9, -10, -11, and -12) cleave execute apoptosis by cleaving targ tory element binding proteins (SRE parane attachment domain. Caspase p17 and p19 fragment of Caspase l length caspase-3. The cleaved P1 n blot (PMID: 25501826). bmed ID Journal 939534 J Membr Biol 798643 Arch Biochem Bio 817595 CNS Neurosci Th er shipment.	and activate downstream effector caspases eted cellular proteins. Caspase 3 plays a key EBPs) between the basic helix-loop-helix 3 can form heterocomplex with other 3. This antibody is specific for cleaved 7 and P19 fragment might form complex and <b>Application</b> WB ophys WB,IF	
	Caspases, a family of endoproteases, cell death. Initiator caspases (caspase (caspase-3, -6, and -7), which in turn errole in the activation of sterol regular leucine zipper domain and the memb proteins. This antibody can recognize caspase 3, and does not recognize ful shows at around 30-35 kDa by wester Author Pul Fei Hu 399 Bin Xia 399 Qingsheng Meng 399 Storage: Store at -20°C. Stable for one year aft Storage Buffer:	e-2, -8, -9, -10, -11, and -12) cleave execute apoptosis by cleaving targ tory element binding proteins (SRE para attachment domain. Caspase p17 and p19 fragment of Caspase l length caspase-3. The cleaved P1 n blot (PMID: 25501826). bmed ID Journal 939534 J Membr Biol 798643 Arch Biochem Bio 817595 CNS Neurosci Th er shipment.	and activate downstream effector caspases eted cellular proteins. Caspase 3 plays a key EBPs) between the basic helix-loop-helix 3 can form heterocomplex with other 3. This antibody is specific for cleaved 7 and P19 fragment might form complex and <b>Application</b> WB ophys WB,IF	

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82707-13-RR (Cleaved Caspase 3/P17/P19 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.