For Research Use Only

YY1 Recombinant antibody, PBS Only

Catalog Number:82712-3-PBS



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

1B5

82712-3-PBS

BC037308 GeneID (NCBI): Protein A purification

CloneNo.:

100ug, Concentration: 1mg/ml by Nanodrop:

UNIPROT ID:

P25490

Rabbit Isotype:

Full Name: YY1 transcription factor

IgG

Calculated MW:

Immunogen Catalog Number: AG17792

414 aa, 45 kDa Observed MW:

65-70 kDa

Applications

Tested Applications:

WB, IF/ICC, ChIP, Indirect ELISA

Species Specificity:

human, mouse

Background Information

YY1, also named as DELTA, INO 80S and NF-E1, contains four C2H2-type zinc fingers and belongs to the YY transcription factor family. YY1 is a multifunctional transcription factor that exhibits positive and negative control on a large number of cellular and viral genes by binding to sites overlapping the transcription start site. YY1 may direct histone deacetylases and histone acetyltransferases to a promoter in order to activate or repress the promoter, thus implicating histone modification in the YY1. The open reading frame of the human YY1 cDNA encodes a protein of 414 amino acids with a predicted molecular weight of 44 kDa. However, YY1 migrates on SDS gels as a 65-68 kDa protein, probably due to the structure of the protein. It is a ubiquitously expressed transcription factor with fundamental roles in embryogenesis, differentiation, replication and proliferation.


Storage

Storage:


Store at -80°C. Storage Buffer:

PBS only, pH7.3


Selected Validation Data




Chromatin was prepared from K-562 cells, cells were fixed with formaldehyde for 10 minutes. The ChIP was performed with 15 μg of cross-linked chromatin, 5 μg of YY1 Antibody (82712-3-RR) or 5 ug of Normal Rabbit IgG (98136-1-RR), and 20 μl of Protein A Magarose Beads. The immunoprecipitated DNA was quantified by real time PCR This data was developed using the same antibody clone with 82712-3-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 82712-3-RR (YY1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 82712-3-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using YY1 antibody (82712-3-RR, Clone: 1B5) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), Alpha Tubulin antibody (66031-1-lg, Clone: 1E4C11, red). This data was developed using the same antibody clone with 82712-3-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using YY1 antibody (82712-3-RR, Clone: 1B5) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 82712-3-PBS in a different storage buffer formulation



Biolayer interferometry (BLL) kinetic assays of 82712-3-RR against Human YY1 were performed. The affinity constant is 0.835 nM.