For Research Use Only

Phospho-MYL9 (Thr19/Ser20) Recombinant antibody

Catalog Number:82717-1-RR



Purification Method:

Basic Information

Catalog Number: GenBank Accession Number:

82717-1-RR BC002648 Protein A purification GeneID (NCBI): CloneNo.: 100ul , Concentration: 1000 ug/ml by 10398 1G10

Nanodrop: **UNIPROT ID:** Recommended Dilutions: Source: P24844 WB 1:5000-1:50000 Rabbit

Full Name: Isotype: myosin, light chain 9, regulatory

IgG Calculated MW:

20 kDa Observed MW: 19-20 kDa

Applications

Tested Applications:

WB, ELISA WB: Calyculin A treated NIH/3T3 cells, Calyculin A

Positive Controls:

Species Specificity: treated HeLa cells human, mouse

Background Information

 $Myosin\ regulatory\ light\ polypeptide\ 9\ (MYL9), also\ known\ as\ MLC2, belongs\ to\ the\ myosin\ regulatory\ subunits.\ It$ plays an important role in regulation of both smooth muscle and nonmuscle cell contractile activity via its phosphorylation at Thr19 and Ser20. Implicated in cytokinesis, receptor capping, and cell locomotion (PMID:11942626, PMID:2526655). Some studies have demonstrated that MYL9 may play important roles in various human cancers. The expression and phosphorylation of MYL9 (Thr19/Ser20) may be increased in human breast (PMID: 22144583) and liver cancers (PMID: 18648664), while decreased in human colon (PMID: 22752057) and bladder cancers (PMID: 21139803). MYL9 was the only gene differentially expressed in the aged versus young injured arteries in the rat smooth muscle cell layers (PMID:22003410).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

Selected Validation Data



Non-treated and Calyculin A treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 82717-1-RR (Phospho-MYL9 (Thr19/Ser20) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Beta Actin antibody as loading control.



1x10^6 Calyculin A treated HeLa cells were intracellularly stained with 0.25 ug Phospho-MYL9 (Thr19/Ser20) Recombinant antibody (82717-1-RR, Clone:1G10) and Coralite® 488-Conjugated Goat Anti-Rabbit 1gG (H+L) (SA00013-2)(red), or 0.25 ug Rabbit 1gG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.