For Research Use Only

## PD-L1/CD274 Recombinant antibody

Catalog Number:82719-13-RR




| Basic Information               | Catalog Number:<br>82719-13-RR   | GenBank Accession Number:<br>BC074984  | Purification Method:<br>Protein A purification                |              |   |                                 |             |  |
|---------------------------------|--|--|---|--------------|---|---------------------------------|-------------|--|
|                                 | Size:<br>100ul , Concentration: 414 ug/ml by<br>Nanodrop;<br>Source:<br>Rabbit<br>Isotype:<br>IgG<br>Immunogen Catalog Number:<br>AG12432  | GenelD (NCBI):<br>29126<br>UNIPROT ID:<br>Q9NZQ7<br>Full Name:<br>CD274 molecule<br>Calculated MW:<br>290 aa, 33 kDa | CloneNo.:<br>3G4<br>Recommended Dilutions:<br>IHC 1:200-1:800 |              |   |                                 |             |  |
|                                 |  |  |   | Applications | Tested Applications:<br>IHC, EUSA   |                                 | e Controls: |  |
|                                 |  |  |   |              | Species Specificity:<br>human   | IHC : human tonsillitis tissue, |             |  |
|                                 |  |  |   |              | Note-IHC: suggested antigen r<br>TE buffer pH 9.0; (*) Alternati<br>retrieval may be performed w<br>buffer pH 6.0 | vely, antigen                   |             |  |
| Background Information          | Programmed cell death ligand 1 (PD-L1, CD274, or B7-H1), is the first member of B7 family to be discovered. B7 family molecules are type I transmembrane proteins belonging to the immunoglobulin superfamily. In concert with their CD28 family receptors, the B7s are key regulators of the adaptive immune response. PD-L1 is suggested as a negative regulator of T and B cell, and plays important role in mediating tolerance of lymphocytes to self-antigens. It is also involved in the costimulatory signal, essential for T-cell proliferation and production of IL10 and IFNG, in an IL2-dependent and a PD-1-independent manner. PD-L1 is a 290 aa transmembrane protein with a calculated molecular weight of 33 kDa, it is predicted to be 27-30 kDa after signal peptide cleavage (PMID: 25609200; 17076679). The apparent molecular weight has also been reported as 45-70 kDa, major glycosylated form of 45-50 kDa and multiple post-translational modifications form of 65-70 kDa (PMID: 18760278; 16493058). |  |   |              |   |                                 |             |  |
| Storage                         | Storage:<br>Store at -20°C. Stable for one year aft<br>Storage Buffer:<br>PBS with 0.02% sodium azide and 50   |  |   |              |   |                                 |             |  |
| *** 20ul sizes contain 0.1% BSA | Aliquoting is unnecessary for -20 $^{\circ}$ C s   | storage  |   |              |   |                                 |             |  |

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 82719-13-RR (PD-L1/CD274 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 82719-13-RR (PD-L1/CD274 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLL) kinetic assays of 82719-13-RR against Human PD-L1/CD274 were performed. The affinity constant is below 1 pM.