

For Research Use Only

CD79A Recombinant antibody

Catalog Number: 82747-4-RR



Basic Information

Catalog Number: 82747-4-RR	GenBank Accession Number: BC113733	Purification Method: Protein A purification
Size: 100ul , Concentration: 600 µg/ml by Nanodrop;	GeneID (NCBI): 973	CloneNo.: 1F24
Source: Rabbit	Full Name: CD79a molecule, immunoglobulin-associated alpha	Recommended Dilutions: WB 1:2000-1:12000
Isotype: IgG	Calculated MW: 226 aa, 25 kDa	
Immunogen Catalog Number: AG17924	Observed MW: 40 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : Daudi cells, Raji cells, Ramos cells
Species Specificity: Human	

Background Information

CD79A, also named as B-cell antigen receptor complex-associated protein alpha chain or MB-1 membrane glycoprotein, is a 226 amino acid protein, which contains one ITAM domain and one Ig-like C2-type (immunoglobulin-like) domain. CD79A is expressed in B cell and localizes in the cell membrane. CD79A is required in cooperation with CD79B for initiation of the signal transduction cascade activated by binding of antigen to the B-cell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. CD79A is also required for BCR surface expression and for efficient differentiation of pro- and pre-B-cells.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82747-4-RR (CD79A antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.