

For Research Use Only

Thioredoxin Recombinant antibody

Catalog Number: 82765-1-RR



Basic Information

Catalog Number: 82765-1-RR	GenBank Accession Number: BC003377	Purification Method: Protein A purification
Size: 100ul , Concentration: 500 µg/ml by Nanodrop;	GeneID (NCBI): 7295	CloneNo.: 7G10
Source: Rabbit	Full Name: thioredoxin	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG	Calculated MW: 12 kDa	
Immunogen Catalog Number: AG6989	Observed MW: 12 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : A549 cells, HeLa cells, HepG2 cells, K-562 cells, MCF-7 cells
Species Specificity: Human	

Background Information

TXN, TRDX, TRX, TRX1, ADF and SASP, belongs to the thioredoxin family. It participates in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyzes dithiol-disulfide exchange reactions. TXN plays a role in the reversible S-nitrosylation of cysteine residues in target proteins, and thereby contributes to the response to intracellular nitric oxide. Nitrosylates the active site Cys of CASP3 in response to nitric oxide (NO), and thereby inhibits caspase-3 activity. TXN induces the FOS/JUN AP-1 DNA-binding activity in ionizing radiation (IR) cells through its oxidation/reduction status and stimulates AP-1 transcriptional activity.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82765-1-RR (Thioredoxin antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.