## For Research Use Only

## Thioredoxin Recombinant antibody

Catalog Number:82765-1-RR



**Basic Information** 

Catalog Number: GenBank Accession Number:

82765-1-RR BC003377 GeneID (NCBI):

100ul , Concentration: 500 ug/ml by Nanodrop: **UNIPROT ID:** P10599 Rabbit Full Name: Isotype: thioredoxin IgG Calculated MW: Immunogen Catalog Number: 12 kDa AG6989

Observed MW: 12 kDa

**Applications** 

**Tested Applications:** WB, IHC, ELISA Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Positive Controls:** 

WB: A549 cells, HeLa cells, HepG2 cells, K-562 cells,

**Purification Method:** 

CloneNo.:

7G10

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000

IHC 1:1000-1:4000

MCF-7 cells

IHC: human stomach tissue,

## **Background Information**

TXN, TRDX, TRX1, ADF and SASP, belongs to the thioredoxin family. It participates in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyzes dithiol-disulfide exchange reactions. TXN plays a role in the reversible S-nitrosylation of cysteine residues in target proteins, and thereby contributes to the response to intracellular nitric oxide. Nitrosylates the active site Cys of CASP3 in response to nitric oxide (NO), and thereby inhibits caspase-3 activity. TXN induces the FOS/JUN AP-1 DNA-binding activity in ionizing radiation (IR) cells through its oxidation/reduction status and stimulates AP-1 transcriptional activity.

Storage


Store at -20°C. Stable for one year after shipment. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 82765-1-RR (Thioredoxin antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human stomach tissue slide using 82765-1-RR (Thioredoxin antibody) at dilution of 1:2000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 82765-1-RR against Human Thioredoxin were performed. The affinity constant is 0.437 nM.