

SIRT1 Recombinant antibody

Catalog Number: 82790-2-RR

Basic Information

Catalog Number: 82790-2-RR	GenBank Accession Number: BC012499	Purification Method: Protein A purification
Size: 100ul , Concentration: 400 µg/ml by Nanodrop;	GeneID (NCBI): 23411	CloneNo.: 4F14
Source: Rabbit	UNIPROT ID: Q96EB6	Recommended Dilutions: WB 1:1000-1:8000
Isotype: IgG	Full Name: sirtuin (silent mating type information regulation 2 homolog) 1 (S. cerevisiae)	
Immunogen Catalog Number: AG3808	Calculated MW: 747 aa, 82 kDa	
	Observed MW: 110-130 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HEK-293 cells, HeLa cells, MCF-7 cells, Jurkat cells, K-562 cells
Species Specificity: Human	

Background Information

SIRT1, also named as SIR2L1, contains a deacetylase sirtuin-type domain and belongs to the sirtuin family. The post-translation modified SIRT1 is a 110-130 kDa protein, which contains one deacetylase sirtuin-type domain. The 75-80 kDa SirT1 fragment was detected to lack the carboxy-terminus (PMID:21305533). SirT1 exists a 57-61 kDa isoform. SIRT1 may be found in nucleolus, nuclear euchromatin, heterochromatin and inner membrane. It can shuttle between nucleus and cytoplasm. SIRT1 regulates processes such as apoptosis and muscle differentiation by deacetylating key proteins. SIRT1 in particular initiates several signaling events relevant to cardioprotection, including: activation of endothelial nitric oxide synthase, INS receptor signaling, and autophagy. In addition SIRT1 activation elicits resistance to oxidative stress via regulation of transcription factors and co-activators such as FOXO, Hif-2a, and NF-κB. SIRT1 regulates the p53-dependent DNA damage response pathway by binding to and deacetylating p53, specifically at Lysine 382.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82790-2-RR (SIRT1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.