

For Research Use Only

MYB/c-Myb Recombinant antibody

Catalog Number: 82804-2-RR



Basic Information

Catalog Number: 82804-2-RR	GenBank Accession Number: BC064955	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 4602	CloneNo.: 230179C11
Source: Rabbit	UNIPROT ID: P10242	Recommended Dilutions: WB 1:2000-1:10000
Isotype: IgG	Full Name: v-myb myeloblastosis viral oncogene homolog (avian)	
Immunogen Catalog Number: AG12021	Calculated MW: 640 aa, 72 kDa	
	Observed MW: 40-50 kDa, 72-80 kDa	

Applications

Tested Applications: WB, FC (Intra), ELISA	Positive Controls: WB : HEK-293 cells, Jurkat cells, MOLT-4 cells, HT-29 cells
Species Specificity: human	

Background Information

The MYB protein is a transcription factor that plays a key role in regulating the proliferation/differentiation of progenitor cells in bone marrow, colonic crypts, and neurogenic niches [PMID:18574464]. It is involved in the regulation of epithelial-to-mesenchymal transition (EMT) and invasion in neuroblastoma, colon carcinoma, and embryonic kidney cells through the upregulation of the transcription repressor Slug [PMID:20622260]. It also a DNA-binding protein that specifically recognize the sequence 5'-YAAC[GT]G-3', and plays an important role in the control of proliferation and differentiation of hematopoietic progenitor cells. MYB exists some isoforms with MV 85, 75, 72, 68, 63 and 40 kDa.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82804-2-RR (MYB antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



1x10⁶ MCF-7 cells were intracellularly stained with 0.25 ug MYB/c-Myb Recombinant antibody (82804-2-RR, Clone:230179C11) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2) (red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).