For Research Use Only

Calretinin Recombinant antibody

Catalog Number:82811-1-RR



Basic Information

Catalog Number: GenBank Accession Number:

82811-1-RR BC015484 Protein A purification GeneID (NCBI): CloneNo.:

29 kDa

100ul , Concentration: 250 ug/ml by 2F20

Nanodrop: **UNIPROT ID:** Recommended Dilutions: P22676 WB 1:1000-1:5100 Rabbit IHC 1:50-1:500 Full Name: IF/ICC 1:125-1:500 Isotype: calbindin 2 IgG Calculated MW:

AG2924 Observed MW:

29 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Immunogen Catalog Number:

Species Specificity: human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

WB: SH-SY5Y cells, U2OS cells, fetal human brain tissue, mouse brain tissue, rat brain tissue, pig brain

Purification Method:

IHC: mouse brain tissue, IF/ICC: SH-SY5Y cells.

Background Information

Calbindin 2 (calretinin), is an intracellular calcium-binding protein belonging to the troponin C superfamily. Members of this protein family have six EF-hand domains which bind calcium. This protein plays a role in diverse cellular functions, including message targeting and intracellular calcium buffering. It also functions as a modulator of neuronal excitability, and is a diagnostic marker for some human diseases, including Hirschsprung disease and some cancers. Calretinin is a useful marker for differentiating malignant mesothelioma from carcinomas.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed adult mouse cerebellum tissue using Calretinin antibody (82811-1-RR) at dilution of 1:500 and Alexa Fluor 594 Affini Pure Donkey Anti-Rabbit IgG (H+L). (Image produced by Watt Lab at McGill University)



Various lysates were subjected to SDS PAGE followed by western blot with 82811-1-RR (Calretinin antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 82811-1-RR (Calretinin antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 82811-1-RR (Calretinin antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed SH-SY5Y cells using Calretinin antibody (82811-1-RR, Clone: 2F20) at dilution of 1:250 and Multi-rAb CoraLite ® Plus 594-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR004).



1x10^6 SH-SY5Y cells were intracellularly stained with 0.25 ug Calretinin Recombinant antibody (82811-1-RR, Clone:2F20) and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).