

For Research Use Only

CLOCK Recombinant antibody, PBS Only

Catalog Number: 82829-1-PBS



Basic Information

Catalog Number: 82829-1-PBS	GenBank Accession Number: BC041878	Purification Method: Protein A purification
Size: 100ug , Concentration: 1mg/ml by Nanodrop;	GeneID (NCBI): 9575	CloneNo.: 1123
Source: Rabbit	UNIPROT ID: O15516	
Isotype: IgG	Full Name: clock homolog (mouse)	
Immunogen Catalog Number: AG30011	Calculated MW: 846 aa, 95 kDa	
	Observed MW: 110 kDa	

Applications

Tested Applications:
WB, IHC, Indirect ELISA

Species Specificity:
Human, mouse, rat

Background Information

Circadian locomotor output cycles protein kaput (CLOCK), also named as BHLHE8 or KIAA0334, is a 846 amino acid protein, which contains one bHLH domain, one PAC domain, and two PAS domains. CLOCK localizes in the nucleus and cytoplasm. CLOCK is expressed in all tissues examined including spleen, thymus, prostate, testis, ovary, small intestine, colon, leukocytes, heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. ARNTL/2-CLOCK heterodimers activate E-box element (5'-CACGTG-3') transcription of a number of proteins of the circadian clock, such as PER1 and PER2. The calculated molecular weight of CLOCK is 95kDa, we detected a 95-110 kDa protein by western blot. Our result is similar to the result of research paper (PMID: 30683868).

Storage


Storage:
Store at -80°C.

Storage Buffer:
PBS Only


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82829-1-RR (CLOCK antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 82829-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 82829-1-RR (CLOCK antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 82829-1-PBS in a different storage buffer formulation.



WB result of CLOCK antibody (82829-1-RR; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CLOCK transfected HeLa cells. This data was developed using the same antibody clone with 82829-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 82829-1-RR (CLOCK antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 82829-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 82829-1-RR (CLOCK antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 82829-1-PBS in a different storage buffer formulation.