For Research Use Only

TAF15 Recombinant antibody

Catalog Number:82860-1-RR



Purification Method:

Basic Information

Catalog Number: GenBank Accession Number:

82860-1-RR BC046099 Protein A purification

GeneID (NCBI): CloneNo.: 100ul, Concentration: 800 µg/ml by 8148 3L13

factor, 68kDa

Recommended Dilutions: Source: TAF15 RNA polymerase II, TATA box WB 1:2000-1:16000 Rabbit binding protein (TBP)-associated IF 1:400-1:1600

Isotype: Calculated MW: IgG 592 aa, 62 kDa Immunogen Catalog Number: AG20828 Observed MW: 62-75 kDa

Applications

Positive Controls: Tested Applications:

IF, WB, ELISA WB: HeLa cells, mouse liver tissue, HEK-293 cells, Species Specificity: Jurkat cells, MCF-7 cells, rat brain tissue

Human, mouse, rat

Background Information

TAF15 (TAF2N or TAFII68) is a nuclear protein known to associate with a distinct subpopulation of transcription factor IID (TFIID), a multi-subunit complex that nucleates the pre-initiation complex on promoters of on proteincoding genes. TAF15 harbors a transcriptional activation domain, a RNA recognition motif and many Arg-Gly-Gly repeats that participate in RNA binding. TAF15 along with FUS and EWS constitutes the FET (FUS/EWS/TAF15) protein family of heterogeneous nuclear ribonucleoproteins (hnRNPs) class of RNA binding proteins that are multifunctional proteins implicated in numerous aspects of RNA processing/functions. TAF15 is a nuclear protein that shuttle between cytoplasm-nucleus and as RNA/ssDNA-binding protein, it play specific roles during transcription initiation at distinct promoters and enter the preinitiation complex together with the RNA polymerase II. TAF15 as well as other FET member genes are frequently translocated in sarcomas and rare hematopoietic as $well \ as \ epithelial \ cancers. \ Moreover, dominant \ mutations \ in \ genes \ coding \ for \ four \ hnRNPs \ (TDP-43, FUS/TLS, TAF15, TA$ and EWS) are found in familial and sporadic cases of ALS and a chromosomal aberration involving TAF15/TAF2N is found in a form of extraskeletal myxoid chondrosarcomas (EMC).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82860-1-RR (TAF15 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 82860-1-RR (TAF15 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using TAF15 antibody (82860-1-RR, Clone: 3L13) at dilution of 1:800 and Multi-rAb CoraLite ® Plus 594-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR004).CoraLite® Plus 488-Phalloidin (green)