## For Research Use Only

## NOP58 Recombinant antibody

Catalog Number:82879-1-RR **Featured Product** 



**Basic Information** 

Catalog Number: GenBank Accession Number:

82879-1-RR BC032592

GeneID (NCBI): Size: 100ul, Concentration: 1000 ug/ml by 51602

Nanodrop: **UNIPROT ID:** Q9Y2X3

Rabbit Full Name: Isotype: NOP58 ribonucleoprotein homolog

IgG (veast)

Immunogen Catalog Number: Calculated MW:

60 kDa AG5740

> Observed MW: 60 kDa

**Purification Method:** 

Protein A purification

CloneNo.: 230058C7

Recommended Dilutions:

WB 1:2000-1:10000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

IHC 1:500-1:2000 IF/ICC 1:500-1:2000

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HeLa cells, MCF-7 cells, Jurkat cells, K-562 cells,

HEK-293 cells IP: HeLa cells,

IHC: human prostate cancer tissue,

IF/ICC: U2OS cells,

## **Background Information**

The box C/D snoRNPs are named for the short, conserved box C (UGAUGA) and box D (CUGA) sequences present in their RNA moiety. The box C/D sequences are also required for multiple aspects. of snoRNA biogenesis, including RNA stability, intronic processing, nucleolar targeting, nuclear retention, and 59 trimethylguanosine cap formation (PMID:9649444,8878486). Nop58 is a component common to the box C/D small nucleolar ribonucleoprotein (PMID:10606270).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82879-1-RR (NOP58 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




IP result of anti-NOP58 (IP:82879-1-RR, 4ug; Detection:82879-1-RR 1:2000) with HeLa cells lysate 1600 ug.




WB result of NOP58 antibody (82879-1-RR; 1:120000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NOP58 transfected MCF-7 cells.




Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 82879-1-RR (NOP58 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed U2OS cells using NOP58 antibody (82879-1-RR) at dilution of 1:1000 and Multi-rAb CoraLite ® Plus 594-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR004).



1x10^6 HeLa cells were intracellularly stained with 0.25 ug NOP58 Recombinant antibody (82879-1-RR, Clone:230058C7) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2) (red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Biolayer interferometry (BLL) kinetic assays of 82879-1-RR against Human NOP58 were performed. The affinity constant is below 1 pM.



Biolayer interferometry (BLL) kinetic assays of 82879-1-RR against Human NOP58 were performed. The affinity constant is 5.53 nM.