

For Research Use Only

RBM17 Recombinant antibody

Catalog Number: 82889-3-RR

Featured Product



Basic Information

Catalog Number:

82889-3-RR

Size:

100ul, Concentration: 1000 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0985

GenBank Accession Number:

BC007871

GeneID (NCBI):

84991

UNIPROT ID:

Q96I25

Full Name:

RNA binding motif protein 17

Calculated MW:

45 kDa

Observed MW:

45-50 kDa

Purification Method:

Protein A purification

CloneNo.:

230125A3

Recommended Dilutions:

WB 1:2000-1:10000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:300-1:1200

Applications

Tested Applications:

WB, IHC, IP, ELISA

Species Specificity:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : BxPC-3 cells, MCF-7 cells

IP : MCF-7 cells,

IHC : human ovarian cancer,

Background Information

RNA-binding motif protein 17 (RBM17), also named splicing factor 45, was first identified as a member of the spliceosome. RBM17 expression is low in normal tissues but is overexpressed in several forms of cancer, including breast, ovarian and prostate.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82889-3-RR (RBM17 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




WB result of RBM17 antibody (82889-3-RR; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-RBM17 transfected MCF-7 cells.



IP result of anti-RBM17 (IP:82889-3-RR, 4ug; Detection:82889-3-RR 1:10000) with MCF-7 cells lysate 1470 ug.



Immunohistochemical analysis of paraffin-embedded human ovarian cancer slide using 82889-3-RR (RBM17 antibody) at dilution of 1:600 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 82889-3-RR against Human RBM17 were performed. The affinity constant is 0.828 nM.