

For Research Use Only

SLC30A1 Recombinant antibody

Catalog Number: 82895-2-RR



Basic Information

Catalog Number: 82895-2-RR	GenBank Accession Number: BC121015	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 7779	CloneNo.: 230152E10
Source: Rabbit	UNIPROT ID: Q9Y6M5	Recommended Dilutions: WB 1:2000-1:10000
Isotype: IgG	Full Name: solute carrier family 30 (zinc transporter), member 1	
Immunogen Catalog Number: AG18567	Calculated MW: 507 aa, 55 kDa	
	Observed MW: 55-63 kDa	

Applications

Tested Applications: WB, FC, ELISA	Positive Controls: WB : SW480 cells, HeLa cells, HepG2 cells
Species Specificity: Human	

Background Information

Zinc transporter 1 (ZNT1), also known as SLC30A1, is the only zinc transporter predominantly located on the plasma membrane, where it plays a pivotal role exporting cytosolic zinc to the extracellular space. SLC30A1 expression is posttranslationally regulated by cellular zinc status and can be N-glycosylated on Asn299 in the extracellular loop between transmembrane domains (PMID: 15451416, 18936201). Western analysis detected a higher band at 63 kDa than the expected molecular mass, which could be attributed to the properties of the cytosolic histidine-rich loop (PMID: 31471319).

Storage



Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82895-2-RR (SLC30A1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.

1x10⁶ U2OS cells were intracellularly stained with 0.25 ug SLC30A1 Recombinant antibody (82895-2-RR, Clone:230152E10) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).