For Research Use Only

AURKA Recombinant antibody

Catalog Number:82906-2-RR Featured Product




eneID (NCBI): 90 IIPROT ID: 14965 Il Name: rora kinase A Iculated MW:	CloneNo.: 230087F7 Recommended Dilutions: WB 1:5000-1:50000 IF/ICC 1:200-1:800
14965 Il Name: rora kinase A	WB 1:5000-1:50000
kDa oserved MW: -52 kDa	
Positive Co WB : K-562 NCCIT cell	cells, HEK-293 cells, MCF-7 cells, HeLa cells,
IF/ICC : HepG2 cells,	
llation during anaphase and/or t ent evidence revealed that AUR expressed highly in testis and v	he Ser/Thr protein kinase family. This telophase by being involved in microtubule RKA may also be implicated in tumor rarious proliferating cells, migrating as a 46 tent in normal tissue but strong in tumor
g	shipment. glycerol pH 7.3. age


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82906-2-RR (AURKA antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using AURKA antibody (82906-2-RR, Clone: 230087F7) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1).



WB result of AURKA antibody (82906-2-RR; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-AURKA transfected HEK-293 cells.



Biolayer interferometry (BLL) kinetic assays of 82906-2-RR against Human AURKA were performed. The affinity constant is 76.9 pM.