## For Research Use Only

## CD147 Recombinant antibody

Catalog Number:82909-1-RR



**Purification Method:** 

CloneNo.:

Protein A purification

**Basic Information** 

**Applications** 

Catalog Number: GenBank Accession Number:

82909-1-RR BC009040 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 682 230046C1 Nanodrop: **UNIPROT ID:** Recommended Dilutions: Source: P35613 IF/ICC 1:500-1:2000

Rabbit Full Name:

Isotype: basigin (Ok blood group)

IgG Calculated MW: 385 aa, 42 kDa

**Tested Applications:** Positive Controls:

IF/ICC, ELISA IF/ICC: HeLa cells, MCF-7 cells Species Specificity:

**Background Information** 

CD147, also known as Basigin or extracellular matrix metalloproteinase inducer (EMMPRIN), is a transmembrane glycoprotein that belongs to the immunoglobulin superfamily (PMID: 7812975). The molecule is composed of an intracellular portion, an extracellular portion and a single transmembrane region. CD147 is expressed on a variety of cell types (e.g., hematopoietic, epithelial, and endothelial cells) and at varying levels (PMID: 32968061). Increased expression of CD147 occurs in many tumors. CD147 is a pleiotropic molecule that plays an important role in fetal, neuronal, lymphocyte and extracellular matrix development (PMID: 17945211). CD147 has been identified as a receptor essential for erythrocyte invasion by Plasmodium falciparum (PMID: 22080952). It has been reported that spike protein of SARS-CoV-2 binds to CD147 on host cells, thereby mediating the viral invasion (Wang, Ke, et al, BioRxiv. 2020).

Storage Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

human

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

## **Selected Validation Data**



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CD147 antibody (82909-1-RR, Clone: 230046C1) at dilution of 1:1000 and MultirAb Coralite ® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM004).



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using CD147 antibody (82909-1-RR, Clone: 230046C1) at dilution of 1:500 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).