# For Research Use Only

# **TBCD** Recombinant antibody

Catalog Number:82911-1-RR

Featured Product



# **Basic Information**

Catalog Number:GenB82911-1-RRBC00Size:Gene100ul, Concentration: 300 ug/ml by6904Nanodrop;UNIPSource:Q9BTRabbitFull NIsotype:tubulIgGCalculImmunogen Catalog Number:133 kAG6310Obse

GenBank Accession Number: BC003094 GeneID (NCBI): 6904 UNIPROT ID: Q9BTW9 Full Name: tubulin folding cofactor D Calculated MW: 133 kDa Observed MW: 130 kDa

### Purification Method: Protein A purification CloneNo.: 230209H7 Recommended Dilutions: WB 1:500-1:3000

WB 1:500-1:3000 IF/ICC 1:50-1:500

# **Applications**

Tested Applications: WB, IF/ICC, FC (Intra), ELISA Species Specificity: human

#### Positive Controls:

WB : Jurkat cells, MCF-7 cells, HeLa cells, HepG2 cells IF/ICC : MCF-7 cells,

### Storage


\*\*\* 20ul sizes contain 0.1% BSA

Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


# Selected Validation Data




WB result of TBCD antibody (82911-1-RR; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-TBCD transfected MCF-7 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 82911-1-RR (TBCD antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using TBCD antibody (82911-1-RR, Clone: 230209H7) at dilution of 1:150 and MultirAb CoraLite ® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM004).





1x10^6 MCF-7 cells were intracellularly stained with 0.25 ug TBCD Recombinant antibody (82911-1-RR, Clone:230209H7) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).

Biolayer interferometry (BLI) kinetic assays of 82911-1 -RR against Human TBCD were performed. The affinity constant is 1.52 nM.