For Research Use Only

COXIV Recombinant antibody

Catalog Number:82916-1-RR

Featured Product



Basic Information

Catalog Number: GenBank Accession Number:

82916-1-RR BC021236 Protein A purification GeneID (NCBI): Size: CloneNo.:

100ul, Concentration: 1000 ug/ml by 1327 3K20 Nanodrop: **UNIPROT ID:** Recommended Dilutions: WB 1:5000-1:50000

Source: P13073 Rabbit Full Name:

Isotype: cytochrome c oxidase subunit IV isoform 1 IgG

Immunogen Catalog Number: Calculated MW: AG1640 19.6 kDa Observed MW:

17-18 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, HepG2 cells, mouse skeletal muscle tissue, Jurkat cells, A431 cells, rat skeletal muscle tissue, mouse liver tissue

Purification Method:

IHC 1:600-1:2000

IF/ICC 1:200-1:800

IHC: human stomach tissue, human colon tissue, human ovarian cancer

IF/ICC: HepG2 cells,

Background Information

COX4I1, also named as COX4 and COXIV-1, belongs to the cytochrome c oxidase IV family. It is one of the nuclearcoded polypeptide chains of cytochrome c oxidase, the terminal oxidase in mitochondrial electron transport. COX4l1 is a marker for mitochondria. It has two isoforms (isoform 1 and 2). Isoform 1(COX4l1) is ubiquitously expressed and isoform 2 is highly expressed in lung tissues.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




WB result of COXIV antibody (82916-1-RR; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-COXIV transfected HepG2 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 82916-1-RR (COXIV antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human normal stomach slide using 82916-1-RR (COXIV antibody) at dilution of 1:2000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using COXIV antibody (82916-1-RR, Clone: 3K20) at dilution of 1:400 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Rabbit IgG(H+L).




Various lysates were subjected to SDS PAGE followed by western blot with 82916-1-RR (COXIV antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 82916-1-RR (COXIV antibody) at dilution of 1:2000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 HeLa cells were intracellularly stained with 0.25 ug COXIV Recombinant antibody (82916-1-RR, Clone:3K20) and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2) (red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLL) kinetic assays of 82916-1-RR against Human COXIV were performed. The affinity constant is 9.41 nM.