

For Research Use Only

STMN2 Recombinant antibody

Catalog Number: 82925-2-RR



Basic Information

Catalog Number: 82925-2-RR	GenBank Accession Number: BC006302	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 11075	CloneNo.: 230110B12
Source: Rabbit	UNIPROT ID: Q93045	Recommended Dilutions: WB 1:2000-1:10000
Isotype: IgG	Full Name: stathmin-like 2	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
Immunogen Catalog Number: AG0929	Calculated MW: 21 kDa	IHC 1:200-1:800
	Observed MW: 18-21 kDa	IF-P 1:250-1:1000
		IF/ICC 1:50-1:500

Applications

Tested Applications: WB, IHC, IF/ICC, IF-P, FC (Intra), IP, ELISA	Positive Controls: WB : SH-SY5Y cells,
Species Specificity: human, mouse	IP : SH-SY5Y cells,
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0 ; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IHC : mouse brain tissue,
	IF-P : mouse brain tissue,
	IF/ICC : SH-SY5Y cells,

Background Information

Stathmin-2 (encoded by STMN2), a member of the stathmin family of proteins, encodes a phosphoprotein that has an important role in microtubule dynamics. The STMN2 protein promotes microtubule dynamics in axonal growth cones, and is thought to play an important role in neurite outgrowth. STMN2 protein is localized in both Golgi and cytosolic compartments. Its expression appears modulated in osteoblasts by nerve growth factor, dexamethasone or RhoA kinase inhibitor Y-27632 which are known effectors of osteogenesis. (PMID: 33841129, PMID: 18611392)

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




SH-SY5Y cells were subjected to SDS PAGE followed by western blot with 82925-2-RR (STMN2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse brain tissue using STMN2 antibody (82925-2-RR, Clone: 230110B12) at dilution of 1:500 and Multi-rAb CoraLite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Ethanol) fixed SH-SY5Y cells using STMN2 antibody (82925-2-RR, Clone: 230110B12) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 82925-2-RR (STMN2 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-STMN2 (IP:82925-2-RR, 4ug; Detection:82925-2-RR 1:1500) with SH-SY5Y cells lysate 1240 ug.



1x10⁶ SH-SY5Y cells were intracellularly stained with 0 ug STMN2 Recombinant antibody (82925-2-RR, Clone:230110B12) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2) (red), or 0 ug Rabbit IgG control Rabbit PolyAb (30000-O-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLI) kinetic assays of 82925-2-RR against Human STMN2 were performed. The affinity constant is 1.48 nM.