For Research Use Only

PAK7 Recombinant antibody

Catalog Number:82935-1-RR

Featured Product



Basic Information

Catalog Number:

GenBank Accession Number: BC024179

Purification Method:

82935-1-RR

GeneID (NCBI):

Protein A purification

Size:

CloneNo.:

100ul , Concentration: 400 ug/ml by

57144

230201A10

Nanodrop:

UNIPROT ID: Q9P286

Recommended Dilutions: WB 1:500-1:2000

Rabbit

Full Name:

Isotype:

p21 protein (Cdc42/Rac)-activated

IgG

kinase 7

Immunogen Catalog Number: AG29190

Calculated MW: 719 aa, 81 kDa

Observed MW:

70 kDa

Applications

Tested Applications:

Positive Controls:

WB, FC (Intra), ELISA

WB: HeLa cells, SH-SY5Y cells

Species Specificity:

Background Information

PAK7, also known as PAK5, is a newly found but scarcely researched type II PAKs (PAK4, 5, 6). PAK7 was originally found in the brain, which promotes filopodia formation in nerve cells. Recently, studies have found that the expression level of PAK7 significantly increased in colorectal cancer, ovarian cancer, gastric cancer, neuroglioma, hepatocellular carcinoma, pancreatic cancer, osteosarcoma, and breast cancer. PAK7 plays an important role in tumorigenesis, invasion, and metastasis, mainly related to its mechanism of regulating cytoskeleton, inhibiting apoptosis, and promoting proliferation. The calculated molecular weight of PAK7 is 80 kDa (PMID: 29805709, PMID: 34165002).

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



WB result of PAK7 antibody (82935-1-RR; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PAK7 transfected HeLa cells.



Various lysates were subjected to SDS PAGE followed by western blot with 82935-1-RR (PAK7 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



(blue)1x10^6 K-562 cells were intracellularly stained with 0.25 ug Anti-Human PAK7 (82935-1-RR, Clone:230201A10) and APC-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) at dilution 1:166(red), or 0.25 ug Rabbit 1gG control Rabbit PolyAb (30000-0-AP). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).