

For Research Use Only

IFT140 Recombinant antibody

Catalog Number: 82947-1-RR



Basic Information

Catalog Number: 82947-1-RR	GenBank Accession Number: BC035577	Purification Method: Protein A purification
Size: 100ul , Concentration: 300 µg/ml by Nanodrop;	GeneID (NCBI): 9742	CloneNo.: 230231F7
Source: Rabbit	UNIPROT ID: Q96RY7	Recommended Dilutions: WB 1:1000-1:6000
Isotype: IgG	Full Name: intraflagellar transport 140 homolog (Chlamydomonas)	
Immunogen Catalog Number: AG11119	Calculated MW: 1462 aa, 165 kDa	
	Observed MW: 150-165 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HeLa cells, rat testis tissue
Species Specificity: Human, Rat	

Background Information

IFT140 is a subunit of intraflagellar transport complex A (IFT-A) which is involved in retrograde ciliary transport. RT-PCR analysis showed it is highly expressed in kidney, moderately in ovary, testis, prostate, and lung. IFT140 is localized to the base and tip of primary cilium. IFT140 has a pivotal role in development and function of ciliated cells, and mutations of IFT140 cause skeletal, renal, and retinal ciliopathies. It had been detected as a single band around 140-165 kDa in different reports. (PMID: 20368623, 22282595)

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 82947-1-RR (IFT140 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



rat testis tissue were subjected to SDS PAGE followed by western blot with 82947-1-RR (IFT140 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.