For Research Use Only

N-cadherin Recombinant antibody

Catalog Number:82968-1-RR



Basic Information

Catalog Number: GenBank Accession Number:

82968-1-RR NM 001792.4 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 1000 Nanodrop:

UNIPROT ID: Source: P19022-1 Rabbit Full Name:

Isotype: cadherin 2, type 1, N-cadherin IgG

(neuronal) Calculated MW:

100 kDa Observed MW: 130 kDa

Purification Method:

Protein A purification

CloneNo.: 230259H5

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:50-1:500 IF/ICC 1:125-1:500

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA

Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, HEK-293 cells

IHC: human intrahepatic cholangiocarcinoma tissue,

human ovary cancer tissue

IF/ICC: HeLa cells,

Background Information

 $Cadherins\ are\ a\ family\ of\ transmembrane\ gly coproteins\ that\ mediate\ calcium-dependent\ cell-cell\ adhesion\ and$ play an important role in the maintenance of normal tissue architecture. N-cadherin (neural cadherin), also known as CDH2 (cadherin 2), is a 130-kDa transmembrane protein and a classical member of the cadherin superfamily which also includes E-, P-, R-, and B-cadherins. Expression of N-cadherin has been reported on various cell types including neurons, endothelial cells and cardiac myocytes (PMID: 11282032; 9508779; 8125202). N-cadherin has functions in early brain morphogenesis, synaptogenesis and synaptic plasticity (PMID: 23321619). The N-cadherin ectodomain can be cleaved, leading to the generation of a 90 kD N-terminal fragment (PMID: 16998833; 17028923).

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82968-1-RR (N-cadherin antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human intrahepatic cholangiocarcinoma tissue slide using 82968-1-RR (N-cadherin antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using N-cadherin antibody (82968-1-RR, Clone: 230259H5) at dilution of 1:250 and CoraLite®488-Conjugated Affini Pure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Immunofluorescent analysis of (-20°C Methanol) fixed A 549 cells using N-cadherin antibody (82968-1-RR, Clone: 230259H5) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).



Biolayer interferometry (BLL) kinetic assays of 82968-1-RR against Human N-cadherin were performed. The affinity constant is 0.464 nM.