For Research Use Only

GDNF Recombinant antibody

Catalog Number:83087-1-RR



Basic Information

Catalog Number: GenBank Accession Number: **Purification Method:**

83087-1-RR BC069369 Protein A purification GeneID (NCBI): CloneNo.:

100ul, Concentration: 1000 ug/ml by 2668 230358A10 Nanodrop:

UNIPROT ID: Recommended Dilutions: P39905 IHC 1:500-1:2000 Rabbit IF-P 1:125-1:500 Full Name: glial cell derived neurotrophic factor IF/ICC 1:50-1:500 Isotype:

IgG Calculated MW: Immunogen Catalog Number: 211 aa, 24 kDa AG24068 Observed MW: 32-40 kDa

Applications

Tested Applications:

IHC, IF/ICC, IF-P, FC (Intra), ELISA

Species Specificity: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

IHC: mouse brain tissue, IF-P: mouse brain tissue, IF/ICC: SH-SY5Y cells,

Background Information

Glial-derived neurotrophic factor (GDNF) is a member of the transforming growth factor (TGF)-a family and exhibits potent neuroprotective activity. GDNF is a potent promoter of neuronal survival in the CNS and peripheral nervous system (PNS). It has a relatively high speciicity for dopaminergic neurons and thus become an efective treatment for Parkinson's diseases (PD), with clinical trials currently in progress. Protein dimers of GDNF have a reported molecular weight of 32-40 kDa(PMID: 17982417, PMID: 23251488, PMID: 11429269).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 83087-1-RR (GDNF antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 83087-1-RR (GDNF antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




(blue)1x10^6 U-251 cells were intracellularly stained with 0.25 ug Anti-Human GDNF (83087-1-RR, Clone:230358A10) and APC-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) at dilution 1:166(red), or 0.25 ug Rabbit 1gG control Rabbit PolyAb (30000-0-AP). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).




1x10^6 SH-SY5Y cells were intracellularly stained with 0.25 ug Anti-Human GDNF (83087-1-RR, Clone:230358A10) and APC-Conjugated Affini Pure Goat Anti-Rabbit 1gG(H-L)(red), or 0.25 ug Rabbit 1gG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse brain tissue using GDNF antibody (83087-1-RR, Clone: 230358A10) at dilution of 1:250 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse brain tissue using GDNF antibody (83087-1-RR, Clone: 230358A10) at dilution of 1:250 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed SH-SY5Y cells using GDNF antibody (83087-1-RR, Clone: 230358A10) at dilution of 1:200 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).