For Research Use Only

TNFR2 Recombinant antibody

Catalog Number:83101-1-RR



Basic Information

Catalog Number:

83101-1-RR

Source

Rabbit

IgG

GenBank Accession Number:

BC052977

GeneID (NCBI):

100ul , Concentration: 1000 $\mu g/ml$ by 7133 Nanodrop:

UNIPROT ID: P20333

Full Name:

Isotype: tumor necrosis factor receptor

superfamily, member 1B

Calculated MW: 48 kDa

Observed MW: 70-75 kDa

Applications

Tested Applications:

WB, ELISA

Species Specificity:

Human

Purification Method: Protein A purification

CloneNo.:

230328B6

Recommended Dilutions:

WB 1:2000-1:10000

Background Information

Tumor necrosis factor-alpha (TNFA/TNFSF2) is a multifunctional cytokine that plays a key role in regulating inflammation, immune functions, host defense, and apoptosis (PMID: 16407280). TNFA signals through two distinctcell surface receptors, TNFR1 (TNFRSF1A, CD120a, p55) and TNFR2 (TNFRSF1B, CD120b, p75). TNFR1 is widely expressed, whereas TNFR2 exhibits more restricted expression, being found on CD4 and CD8 T lymphocytes, endothelial cells, microglia, oligodendrocytes, neuron subtypes, cardiac myocytes, thymocytes and human mesenchymal stem cells (PMID: 20489699; 22374304). In contrast to TNFR1, TNFR2 does not have a death domain. TNFR2 only signals for antiapoptotic reactions. However, recent evidence indicates that TNFR2 also signals to induce TRAF2 degradation (PMID: 22374304). Various defects in the TNFR2 pathway, due to polymorphisms in the TNFR2 gene, upregulated expression of TNFR2 and TNFR2 shedding, have been implicated in the pathology of several autoimmune disorders (PMID: 20489699).

Positive Controls:

WB: NK-92 cells, THP-1 cells

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 98010-2-RR (TNFR2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.