

For Research Use Only

TPR Recombinant antibody

Catalog Number: 83177-5-RR



Basic Information

Catalog Number: 83177-5-RR	GenBank Accession Number: NM_003292	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 7175	CloneNo.: 230413A2
Source: Rabbit	UNIPROT ID: P12270	Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500 IF/ICC 1:50-1:500
Isotype: IgG	Full Name: translocated promoter region (to activated MET oncogene)	
	Calculated MW: 267 kDa	
	Observed MW: ~270 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA	Positive Controls:
Species Specificity: human	WB : HeLa cells, HEK-293 cells
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IHC : mouse testis tissue,
	IF/ICC : HeLa cells,

Background Information

Nucleoporin TPR was originally identified as the oncogenic activator of the met, raf, and trk protooncogenes. TPR is a large 270 kDa protein with a bipartite structure consisting of a ~1600-residue a-helical coiled-coil N-terminal domain and a highly acidic non-coiled ~800 amino acid carboxy terminus (PMID: 22253824). TPR is localized exclusively to intranuclear filaments associated with the nucleoplasmic side of the NPC, by directly binding to Nup153 (PMID: 12802065).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83177-5-RR (TPR antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 83177-5-RR (TPR antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using TPR antibody (83177-5-RR, Clone: 230413A2) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).