

For Research Use Only

HIRA Recombinant antibody, PBS Only

Catalog Number: 83194-4-PBS



Basic Information

Catalog Number: 83194-4-PBS	GenBank Accession Number: NM_003325	Purification Method: Protein A purification
Size: 100ug , Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 7290	CloneNo.: 240006F8
Source: Rabbit	UNIPROT ID: P54198	
Isotype: IgG	Full Name: HIR histone cell cycle regulation defective homolog A (S. cerevisiae)	
Immunogen Catalog Number: AG34439	Calculated MW: 112 kDa	
	Observed MW: 112 kDa	

Applications

Tested Applications:
WB, Indirect ELISA

Species Specificity:
human

Background Information

HIRA protein is encoded by a gene within a region of human chromosome 22q11.2 deleted in most patients with the DiGeorge syndrome, a developmental disorder, and was named for its amino acid sequence homology to two *S. cerevisiae* proteins, Hir1p and Hir2p. HIRA is critical in a specific chromatin assembly pathway that takes place independently of DNA synthesis (PMID: 12049744). HIRA complex associates with various proteins, such as transcription factors, RNA pol II, Prohibitin (PHB), and replication protein A (RPA) suggests that the targeting of the complex at specific locations may exploit additional structural properties and mechanisms (PMID: 30082790).

Storage


Storage:
Store at -80°C.

Storage Buffer:
PBS Only


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83194-4-RR (HIRA antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 83194-4-PBS in a different storage buffer formulation.



1x10⁶ K-562 cells were intracellularly stained with 0.25 ug HIRA Recombinant antibody (83194-4-RR, Clone:240006F8) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 83194-4-PBS in a different storage



Biolayer interferometry (BLI) kinetic assays of 83194-4-RR against Human HIRA were performed. The affinity constant is below 1 pM.