

For Research Use Only

HIRA Recombinant antibody

Catalog Number: 83194-4-RR



Basic Information

Catalog Number: 83194-4-RR	GenBank Accession Number: NM_003325	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 7290	CloneNo.: 240006F8
Source: Rabbit	UNIPROT ID: P54198	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG	Full Name: HIR histone cell cycle regulation defective homolog A (S. cerevisiae)	
Immunogen Catalog Number: AG34439	Calculated MW: 112 kDa	
	Observed MW: 112 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HeLa cells, HEK-293 cells, HepG2 cells, K-562 cells
Species Specificity: human	

Background Information

HIRA protein is encoded by a gene within a region of human chromosome 22q11.2 deleted in most patients with the DiGeorge syndrome, a developmental disorder, and was named for its amino acid sequence homology to two *S. cerevisiae* proteins, Hir1p and Hir2p. HIRA is critical in a specific chromatin assembly pathway that takes place independently of DNA synthesis (PMID: 12049744). HIRA complex associates with various proteins, such as transcription factors, RNA pol II, Prohibitin (PHB), and replication protein A (RPA) suggests that the targeting of the complex at specific locations may exploit additional structural properties and mechanisms (PMID: 30082790).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83194-4-RR (HIRA antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



1×10^6 K-562 cells were intracellularly stained with 0.25 μ g HIRA Recombinant antibody (83194-4-RR, Clone: 240006F8) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2) (red), or 0.25 μ g Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Biolayer interferometry (BLI) kinetic assays of 83194-4-RR against Human HIRA were performed. The affinity constant is below 1 pM.