

For Research Use Only

NDUFB8 Recombinant antibody

Catalog Number: 83216-3-RR



Basic Information

Catalog Number:

83216-3-RR

Size:

100ul, Concentration: 1000 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG6495

GenBank Accession Number:

BC000466

GeneID (NCBI):

4714

UNIPROT ID:

O95169

Full Name:

NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 8, 19kDa

Calculated MW:

22 kDa

Observed MW:

19-22 kDa

Purification Method:

Protein A purification

CloneNo.:

230510B10

Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

IHC 1:50-1:500

IF/ICC 1:125-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HepG2 cells, A549 cells, mouse liver tissue, mouse brain tissue, rat brain tissue, HeLa cells, Jurkat cells

IP : HepG2 cells,

IHC : human stomach cancer tissue, human intrahepatic cholangiocarcinoma tissue

IF/ICC : HepG2 cells,

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83216-3-RR (NDUFB8 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




IP result of anti-NDUFB8 (IP:83216-3-RR, 4ug; Detection:83216-3-RR 1:3000) with HepG2 cells lysate 1470 ug.




Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 83216-3-RR (NDUFB8 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer(pH9).




Immunohistochemical analysis of paraffin-embedded human intrahepatic cholangiocarcinoma tissue slide using 83216-3-RR (NDUFB8 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using NDUFB8 antibody (83216-3-RR, Clone: 230510B10) at dilution of 1:300 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using NDUFB8 antibody (83216-3-RR, Clone: 230510B10) at dilution of 1:250 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



1x10^6 HeLa cells were intracellularly stained with 0.25 ug NDUFB8 Recombinant antibody (83216-3-RR, Clone:230510B10) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLI) kinetic assays of 83216-3-RR against Human NDUFB8 were performed. The affinity constant is 3.26 nM.