

# Nectin-4/PVRL4 Recombinant antibody, PBS Only

Catalog Number: 83223-6-PBS

## Basic Information

<b>Catalog Number:</b> 83223-6-PBS	<b>GenBank Accession Number:</b> BC010423	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 100ug , Concentration: 1 mg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 81607	<b>CloneNo.:</b> 240009E2
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> Q96NY8	
<b>Isotype:</b> IgG	<b>Full Name:</b> poliovirus receptor-related 4	
	<b>Calculated MW:</b> 510 aa, 55 kDa	

## Applications

**Tested Applications:**  
WB, Indirect ELISA

**Species Specificity:**  
Human

## Background Information

Nectin-4, also known as PVRL4, is a type-I transmembrane glycoprotein that belongs to the nectin subfamily of immunoglobulin-like adhesion molecules that participate in Ca(2+)-independent cell-cell adhesion (PMID: 18703497). The extracellular domain of Nectin-4, which contains three immunoglobulin-like domains (V and two C2-type domains, VCC), can be proteolytically cleaved to release a soluble form. Nectin-4 interacts with afadin via its carboxyl-terminal cytoplasmic sequence and trans-interacts with nectin-1/PRR1 through V domain interaction (PMID: 11544254). It acts as a receptor for measles virus (PMID: 22048310). Nectin-4 is overexpressed in multiple human malignancies and the aberrant expression is correlated with cancer progression and poor prognostic (PMID: 35131876). Nectin-4 can be detected as a soluble form of 43 kDa and a transmembrane form of 66 kDa (PMID: 15784625).

## Storage


**Storage:**  
Store at -80°C.

**Storage Buffer:**  
PBS Only

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83223-6-RR (Nectin-4/PVRL4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 83223-6-PBS in a different storage buffer formulation.