For Research Use Only

## gp130/IL6ST Recombinant antibody, PBS Only (Capture)

83284-7-PBS

Size:

Catalog Number:83284-7-PBS



**Purification Method:** 

Protein A purification

CloneNo.:

240992G3

**Basic Information** 

Catalog Number:

GenBank Accession Number:

NM\_002184.3

GeneID (NCBI):

100ug, Concentration: 1 mg/ml by 3572

Nanodrop; UNIPROT ID:
Source: P40189-1
Rabbit Full Name:

Isotype: interleukin 6 signal transducer IgG (gp130, oncostatin M receptor)

Calculated MW: 104 kDa

**Applications** 

**Tested Applications:** 

Sandwich ELISA, Indirect ELISA, Sample test

Species Specificity:

human

**Product Information** 

83284-7-PBS targets gp130/IL6ST as part of a matched antibody pair:

MP00324-3: 83284-7-PBS capture and 83284-4-PBS detection (validated in Sandwich ELISA)


Unconjugated rabbit recombinant monoclonal antibody in PBS only (BSA and azide free) storage buffer at a concentration of 1 mg/mL, ready for conjugation. Created using Proteintech's proprietary in-house recombinant technology. Recombinant production enables unrivalled batch-to-batch consistency, easy scale-up, and future security of supply.

This conjugation ready format makes antibodies ideal for use in many applications including: ELISAs, multiplex assays requiring matched pairs, mass cytometry, and multiplex imaging applications. Antibody use should be optimized by the end user for each application and assay.


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


## **Selected Validation Data**



Sandwich ELISA standard curve of MP00324-3, Human gp130 Recombinant Matched Antibody Pair -PBS only. 83284-7-PBS was coated to a plate as the capture antibody and incubated with serial dilutions of standard Eg0024. 83284-4-PBS was HRP conjugated as the detection antibody. Range: 0.156-10 ng/mL



Serum of sixteen individual healthy human donors was measured. The human gp130 concentration of detected samples was determined to be 123.3 ng/mL with a range of 38.7 - 330.5 ng/mL



Human peripheral blood mononuclear cells (PBMC) (1 x 10^6 cells/mL) were cultured in DMEM supplemented with 8% fetal bonive serum, 5μM β-mercaptoethanol, 2 mM L-glutamine, 100 U/mL penicillin, and 100 μg/mL streptomycin sulfate. Cells were cultured unstimulated or stimulated with 10 μg/mL PHA for 3 days. Aliquots of the cell culture supernates were removed and assayed for levels of human gp130.